



Australian Government




Queensland Government

Marine modelling methods

Reef Water Quality Report Card 2020

Reef 2050 Water Quality Improvement Plan



© State of Queensland, 2022.

The Queensland Government supports and encourages the dissemination and exchange of its information. The copyright in this publication is licensed under a Creative Commons Attribution 4.0 Australia (CC BY) licence.



You must keep intact the copyright notice and attribute the State of Queensland as the source of the publication.

For more information on this licence, visit

<http://creativecommons.org/licenses/by/4.0/au/deed.en>

Disclaimer

This document has been prepared with all due diligence and care, based on the best available information at the time of publication. The government holds no responsibility for any errors or omissions within this document. Any decisions made by other parties based on this document are solely the responsibility of those parties. Information contained in this document is from a number of sources and, as such, does not necessarily represent government or departmental policy.

If you need to access this document in a language other than English, please call the Translating and Interpreting Service (TIS National) on 131 450 and ask them to telephone Library Services on +61 7 3170 5470.

This publication can be made available in an alternative format (e.g. large print or audiotape) on request for people with vision impairment; phone +61 7 3170 5470 or email <library@des.qld.gov.au>.

Citation

Australian and Queensland governments, 2022, *Marine modelling methods, Reef Water Quality Report Card 2020*, State of Queensland, Brisbane.

CONTENTS

- Marine modelling methods..... 4**
- Marine modelling methods..... 5
- eReefs coupled hydrodynamic - biogeochemical model 5
- Modelling improvements 7
- How the metric is calculated and information reported 8
- Qualitative confidence ranking 9
- References 9

Marine modelling methods

The Marine Modelling Program (Waterhouse et al. 2018) directly supports the 2050 outcome of the Reef 2050 Water Quality Improvement Plan (Reef 2050 WQIP) (Australia and Queensland governments), which states:

“Good water quality sustains the outstanding universal value of the Great Barrier Reef, builds resilience, improves ecosystem health and benefits communities.”

The Marine Modelling Program was established in 2016 to:

- Assess trends in ecosystem health for the Great Barrier Reef in relation to water quality and its linkages to end-of-catchment loads by predicting, assessing and reporting trends in inshore water clarity and concentrations of chlorophyll *a*.
- Predict physical and biogeochemical properties of Reef waters under a range of scenarios to assess the impact of management practices and contribute to the establishment or review of basin-level water quality targets.
- Support regional and whole-of-Great Barrier Reef water quality risk assessments by predicting the impact of rivers on the Great Barrier Reef waters under a range of conditions.

Given the scale of the Great Barrier Reef, it would be impractical to measure and report water quality through the entire area and at a reasonable frequency using monitoring data alone.

Satellite imaging can be used to cover this wide spatial area but is generally considered to have lower accuracy and is seasonally affected by cloud cover. Therefore, the eReefs modelling framework is used in conjunction with the monitored information and satellite observations to extrapolate water quality across the entire Great Barrier Reef. The model is used to generate the marine water quality metric.

This report describes the methods used to generate the marine water quality metric for the period 1 October 2019 to 30 September 2020 (water year 2019-2020).

Marine modelling methods

Marine models which integrate physical processes and ecosystem responses play an integral part in supporting resilience-based management and linking science and observations to policy and decision making.

In this context, the marine component of eReefs delivers and operates numerical models capable of simulating and predicting the physical hydrodynamic state, sediment transport, water quality and basal ecology of the Great Barrier Reef lagoon <<https://research.csiro.au/ereefs/models/>>. Together, these models provide the ability to simulate the transport and fate of waterborne material, from the ocean or land, and assess its impact on Reef water quality (Skerratt et al., 2019a).

In 2015-2016, as part of [National Environmental Science Programme \(NESP\) Project 3.2.5](#), eReefs models were used for the first time to report on chlorophyll *a* (productivity linked to nutrient concentrations) and Secchi depth (proxy for water clarity and presence of fine sediments) across the entire Great Barrier Reef (Robillot et al., 2018). These measures underpinned a new water quality metric for the Reef water quality report cards. The metric considered all six regions in calculating the Reef-wide score and is based on open coastal waters.

The new metric is underpinned by the eReefs biogeochemical model and integrates true-colour data from satellite images for improved accuracy in what is commonly referred to as data assimilation (Baird et al., 2016). This integration of multiple streams of data to measure and report on water quality differs from the previous metric, which relied exclusively on model predictions of water quality variables like chlorophyll and suspended sediments derived from satellite data.

The eReefs model has been assessed extensively against *in situ* observations with detailed assessment findings available in the Technical assessment of the eReefs biogeochemical simulation [gbr4_H2p0_B3p0_Chyd_Dcrt] against observations (Skerratt et al., 2019b). The approach to calculating Reef water quality indices and overall scores was independently peer-reviewed as part of NESP Project 3.2.5.

eReefs coupled hydrodynamic - biogeochemical model

The eReefs coupled hydrodynamic, sediment and biogeochemical modelling system involves the application of a range of physical, chemical and biological process descriptions to quantify the rate of change of physical and biological variables.

The process descriptions are generally based either on a fundamental understanding of processes or on actual measurements when a specific process was able to be isolated and studied. The model also requires external inputs, such as observed river flows and pollutant loads.

The three components of the model are:

- The hydrodynamic 3-D model as defined by Herzfeld (2006, 2015).
- The sediment transport model, which adds a multilayer sediment bed to the hydrodynamic model grid and simulates sinking, deposition and resuspension of multiple size classes of suspended sediment (Margvelashvili, 2009, Margvelashvili et al., 2016).
- The biogeochemical model, which simulates optical, nutrient, plankton, benthic organisms (seagrass, macroalgae and coral), detritus, chemical and sediment dynamics across the whole Great Barrier Reef region, spanning estuarine systems to offshore reefs (Figure 1. Skerratt et al., 2019a).




Figure 1: Conceptual framework of the eReefs coupled hydrodynamic-biogeochemical model. Orange variables are optically active (i.e. either scatter or absorb light), influencing the vertical attenuation of light and the bottom light field. The model is forced by rivers along the Reef with nutrient and sediment loads (Baird et al., 2016) using the Source Catchments model. Source: Skerratt et al., 2019a.

Photosynthetic growth is determined by concentrations of dissolved nutrients (nitrogen and phosphorus) and photosynthetically active radiation. Overall, the model contains 23 optically active constituents (Baird et al., 2016). The biogeochemistry model was updated in 2018 to specifically consider ultrafine sediment particles and their impact on matters such as water clarity.

The model is currently forced with freshwater inputs from rivers along the Great Barrier Reef. River flows for 17 rivers are obtained from the Queensland Department of Natural Resources, Mines and Energy gauging network. Nutrient concentrations flowing in from the ocean are obtained from the CSIRO Atlas of Regional Seas (CARS) 2009 climatology (Ridgway et al., 2002).

River pollutant loads are obtained from the Source Catchments modelling outputs (Ellis and Searle, 2013) up to 30 April 2020. To provide daily time series prediction of pollutant loads past 30 April 2020, pollutant generation models are used that estimate daily loads through varying monthly concentrations. These monthly concentration outputs allow the model predictions to be extended by providing daily rainfall run-off model inputs (i.e. the run-off of the day), without the need to update many thousands of farm scale sub-models.

The eReefs model can be run without using observations from the marine environment, which is referred to as a non-assimilating simulation. However, data assimilation provides the single best estimate of the biogeochemical state of the Reef by combining modelling and observations (Jones et al., 2016), and also improves the skill of the model to make predictions (Skerratt et al., 2019a, b). Data assimilation systems can be thought of as using a model to interpolate between observations.

For the Great Barrier Reef, only remote sensing provides the density of observations required to undertake a large-scale data assimilation (see <[eReefs home page | eReefs](#)>).

For shallow inshore waters, using remote sensing to estimate in-water properties is challenging due to the interactions between chlorophyll *a*, sediment, coloured dissolved organic matter and benthic communities, which all absorb and scatter light in the blue and green wavebands.

Instead of using remote sensing to estimate in-water properties, the water quality metric is based on the optical calculations of the biogeochemical model, which simulates the normalised remote-sensing reflectance.

The data assimilation system uses the mismatch between observed and modelled remote-sensing reflectance to constrain the biogeochemical model (Jones et al., 2016).

A 100 member Ensemble Kalman Filter (EnKF) assimilation system is used informed by observed ocean colour from the NASA MODIS Aqua, VIIRS and ESA Sentinel-3A satellites (Jones et al., 2016). When the ocean colour data is ingested, the model shifts a number of optically-active *in situ* quantities, and in particular phytoplankton numbers, in all ensemble members in a manner that is consistent with the statistical properties of the biogeochemical model.

The estimate of the biogeochemical state by the assimilation system is the mean of the 100 ensemble members. While the model assimilates ocean colour data from satellite, it is assessed against the *in situ* observations of chlorophyll *a* concentration, from which the skill of the system can be quantified.

Modelling improvements

The modelling system was initially developed under the eReefs Project and NESP 3.2.5 Project, and improved for Report Card 2017 and 2018. Significant improvements in Report Card 2020 include:

- Incorporating catchment load estimates using the new SOURCE Catchments code with 2020 vegetation cover.
- Improved pre-processing of river loads before input into the model by 'capping' to prevent large river concentrations due to large loads relative to small flows. Loads that cannot be included at the time predicted by SOURCE Catchments are released at the next flow event: this allows concentrations to be below the cap. This mostly affected the O'Connell River due to large ungauged sections of the catchment.

- SOURCE Catchments total suspended sediments were allocated to FineSed (90%) and Dust (10%), replacing constant concentration for Dust of 20 g/m³ used in previous simulations. This reduced the total riverine suspended solids slightly and increased the fraction of ultrafine sediments during large flows.
- From 1 January 2018, Sentinel-3A atmospherically-corrected remote-sensing reflectance was used for the data assimilation.
- Optical properties of suspended inorganic particles were updated based on analysis in Soja-Wozniak et al. (2020). These new values, obtained at Lucinda Jetty Coastal Laboratory, were more representative of the Great Barrier Reef-wide suspended inorganic particles but less representative of coastal embayments.

The revised model has been assessed over a period of eight years and compared to the previous model configuration to confirm it maintained all the properties and skill required to predict water quality.

How the metric is calculated and information reported

The Reef water quality report card marine water quality metric is calculated as follows:

1. Chlorophyll *a* concentration and Secchi depth data are extracted from the assimilated eReefs biogeochemical model at a 4km spatial resolution and daily temporal resolution (midday snapshot) for the entire Reef.
2. The data is partitioned temporally into water years (from 1 October to 30 September of the reporting year) and spatially into zones representing combinations of regions and cross-shelf water bodies (i.e. open coastal, mid-shelf and offshore waters; defined in GBRMPA, 2010). The enclosed coastal water body is excluded due to limitations associated with the 4 km model resolution near the coastline.
3. The site-level data (4 km x 4 km) for each of the three measures are standardised to indices on a continuous scale of zero (very poor) to 100 (very good). This is done by assessing individual values relative to the appropriate water quality guideline value according to a 'modified amplitude indexation routine' (fsMAMP: base 2 logarithm of the ratio of observed value to threshold).
4. Scores for each parameter are aggregated (averaged) temporally over the water year into annual scores and spatially in the open coastal reporting zone. The resulting scores for chlorophyll *a* and Secchi depth are then averaged to generate a single score for each region.
5. A Reef score is calculated as the weighted (relative areas) average of regional scores.
6. All reported scores are mapped onto a five-point (A–E) colour-coded grading scale (see Table 1).

Table 1: Marine water quality metric score to grade scale

| Grade | Status | Range | Colour |
|-------|-----------|--------|-------------|
| E | Very poor | 0–20 | Red |
| D | Poor | 21–40 | Orange |
| C | Moderate | 41–60 | Yellow |
| B | Good | 61–80 | Light green |
| A | Very good | 81–100 | Dark green |

Qualitative confidence ranking

Data confidence



A multi-criteria analysis was used to qualitatively score the confidence in each indicator used in the report card, from low to high. The approach combined expert opinion and direct measures of error for program components where available. Marine modelling received a three-dot confidence ranking.

References

Baird, ME, Cherukuru, N, Jones, E, Margvelashvili, N, Mongin, M, Oubelkheir, K, Ralph, PJ, Rizwi, F, Robson, BJ, Schroeder, T, Skerratt, J, Steven, ADL, Wild-Allen, KA 2016, 'Remote-sensing reflectance and true colour produced by a coupled hydrodynamic, optical, sediment, biogeochemical model of the Great Barrier Reef, Australia: Comparison with satellite data', *Environmental Modelling and Software*, vol. 78, pp. 79-96.

Ellis, R, Searle, R 2013, 'An integrated water quality modelling framework for reporting on Great Barrier Reef catchments', in J Piantadosi, RS Anderssen and J Boland eds, *MODSIM2013, 20th International Congress on Modelling and Simulation*, Modelling and Simulation Society of Australia and New Zealand, December 2013, pp. 3183-89
<www.mssanz.org.au/modsim2013/L21/ellis.pdf>

Great Barrier Reef Marine Park Authority 2010, *Water Quality Guidelines for the Great Barrier Reef Marine Park*, Great Barrier Reef Marine Park Authority, Townsville, Queensland.

Herzfeld, M 2006, 'An alternative coordinate system for solving finite difference ocean models', *Ocean Modelling*, vol. 14 (3-4), pp. 174-196.

Herzfeld, M 2015, 'Methods for freshwater riverine input into regional ocean models', *Ocean Modelling*, vol. 90, pp. 1-15.

Jones, E, Baird, ME, Mongin, M, Parslow, J, Skerratt, J, Lovell, J, Margvelashvili, NY, Matear, R, Wild-Allen, KA, Robson, BJ, Rizwi, F, Oke, P, King, E, Schroeder, T, Steven, ADL and Taylor, J 2016, 'Use of remote-sensing reflectance to constrain a data assimilating marine biogeochemical model of the Great Barrier Reef', *Biogeosciences*, vol. 13, pp. 6441-6469.

Margvelashvili, NY 2009, 'Stretched Eulerian coordinate model for coastal sediment transport', *Computers and Geosciences*, vol. 35, pp. 1167-1176.

Margvelashvili, NY, Herzfeld, M, Rizwi, F, Mongin, M, Baird, ME, Jones, E, Schaffelke, B, King, E and Schroeder, T 2016, 'Emulator-assisted data assimilation in complex models', *Ocean Dynamics*, vol. 66(9), pp.1109-1124.

Ridgway, KR, Dunn, J and Wilkin, J 2002, 'Ocean Interpolation by Four-Dimensional Weighted Least Squares – Application to the Waters around Australasia', *Journal of Atmospheric and Oceanic Technology*, vol. 19, pp.1357-1375.

Robillot, C, Logan, M, Baird, M, Waterhouse, J, Martin, K and Schaffelke, B 2018, 'Testing and implementation of an improved water quality index for the 2016 and 2017 Great Barrier Reef Report Cards', *Report to the National Environmental Science Programme*, Reef and Rainforest Research Centre Limited, Cairns, pp. 65.

Skerratt, JH, Mongin, M, Baird, ME, Wild-Allen, KA, Baird, ME, Robson, BJ, Schaffelke, B, Davies, CH, Richardson, AJ, Margvelashvili, N, Soja-Wozniak, M, and Steven, ADL 2019a, 'Simulated nutrient and plankton dynamics in the Great Barrier Reef (2011-2016)', *Journal of Marine Systems*, vol. 192, pp. 51-74.

Skerratt, JH, Mongin, M, Baird, ME, Wild-Allen, KA, Robson, BJ, Schaffelke, B, Margvelashvili, N and Soja-Wozniak, M 2019b 'Technical assessment of the eReefs biogeochemical (BGC) simulation [gbr4_H2p0_B3p0_Chgd_Dcrt] against observations' CSIRO, Hobart, 165pp. <https://research.csiro.au/ereefs/models/model-outputs/bgcv3p0/gbr4_h2p0_b3p0_chgd_dcrt_4april19/>

Soja-Wozniak, M., M. Baird, T. Schroeder, Y. Qin, L. Clementson, B. Baker, D. Boadle, V. Brando, A. Steven (2019). Particulate backscattering ratio as an indicator of changing particle composition in coastal waters: Observations from Great Barrier Reef waters. *Journal of Geophysical Research: Oceans*, 124. <https://doi.org/10.1029/2019JC014998>.

Waterhouse, J, Lønborg, C, Logan, M, Petus, C, Tracey, D, Lewis, S, Howley, C, Harper, E, Tonin, H, Skuza, M, Doyle, J, Costello, P, Davidson, J, Gunn, K, Wright, M, Zagorskis, I, Kroon, F, Gruber, R 2018, 'Marine Monitoring Program: Annual report for inshore water quality monitoring 2016-2017', Australian Institute of Marine Science, TropWATER.