



Australian Government




Queensland Government

# Great Barrier Reef Marine Monitoring Program Methods

Reef Water Quality Report Card 2020

Reef 2050 Water Quality Improvement Plan



© Great Barrier Reef Marine Park Authority, 2022

Published by the Australian and Queensland governments

This report is licensed for use under a Creative Commons by Attribution 4.0 International licence with the exception of the Coat of Arms of the Australian Government and the Queensland Government, any other material protected by a trademark, content supplied by third parties and any photographs. For licence conditions see:

<http://creativecommons.org/licences/by/4.0>



**This publication should be cited as:**

Great Barrier Reef Marine Park Authority. 2022, *Marine Monitoring Program Methods. Report for the Reef Water Quality Report Card 2020*, Australian and Queensland governments, Brisbane.

The Great Barrier Reef Marine Park Authority acknowledges the continuing Sea Country management and custodianship of the Great Barrier Reef by Aboriginal and Torres Strait Islander Traditional Owners whose rich cultures, heritage values, enduring connections and shared efforts protect the Reef for future generations.

**DISCLAIMER**

While reasonable efforts have been made to ensure that the contents of this document are factually correct, the Great Barrier Reef Marine Park Authority does not make any representation or give any warranty regarding the accuracy, completeness, currency or suitability for any particular purpose of the information or statements contained in this document. To the extent permitted by law the Great Barrier Reef Marine Park Authority shall not be liable for any loss, damage, cost or expense that may be occasioned directly or indirectly through the use of or reliance on the contents of this document.

Comments and questions regarding this document are welcome and should be addressed to:

Great Barrier Reef Marine Park Authority  
280 Flinders Street Townsville | PO Box 1379 Townsville QLD 4810

[info@gbmpa.gov.au](mailto:info@gbmpa.gov.au)

# Table of contents

Marine monitoring methods .....	4
Seagrass condition .....	5
Coral reef condition .....	6
Assessing status against the objectives.....	8
<i>Improved seagrass condition</i> .....	8
<i>Improved coral condition</i> .....	8
<i>Synthesis and integration of data and information</i> .....	9
Qualitative confidence rankings .....	9
References .....	10
Appendix A: Derivation of confidence ranking.....	12
Glossary.....	14

## List of tables

Table 1: Area of seagrass shallower than 15m in each region within the boundaries of the Great Barrier Reef World Heritage Area .....	8
Table 2: Area of inshore reef in each region within the Marine Park .....	9

## List of figures

Figure 1. Marine Monitoring Program seagrass survey locations .....	5
Figure 2: Marine Monitoring Program coral survey locations .....	7
Figure 3: <i>Qualitative confidence rankings for seagrass and coral scores.</i> .....	9

# Marine monitoring methods

This report summarises the coral and seagrass data and methods used for monitoring and reporting within the Great Barrier Reef Marine Monitoring Program (MMP) managed by the Great Barrier Reef Marine Park Authority and reported in the Reef Water Quality Report Card 2020. Detailed methods are available in the [Marine Monitoring Program annual technical report series](#) that undergo independent peer review before being published in the Great Barrier Reef Marine Park Authority's eLibrary.

The Marine Monitoring Program was established in 2005 and assesses trends in ecosystem health and resilience indicators for the inshore Great Barrier Reef in relation to water quality and its linkages to end-of-catchment pollutant loads. The inshore Marine Monitoring Program has three sub-components:

- Water quality, including pesticides (Waterhouse *et al.* 2021)
- Seagrass condition (McKenzie *et al.* 2021)
- Coral reef condition (Thompson *et al.* 2021).

The Marine Monitoring Program is one line of evidence describing the condition and trend of key coral reef and seagrass meadows used to report progress towards the Reef 2050 Water Quality Improvement Plan (Reef 2050 WQIP) (Australia and Queensland governments 2018) 2025 water quality outcome:

- Good water quality sustains the outstanding universal value of the Great Barrier Reef, builds resilience, improves ecosystem health and benefits communities.

The Marine Monitoring Program objectives are:

- Assess temporal and spatial trends in inshore marine water quality and link pollutant concentrations to end-of-catchment loads
- Monitor, assess and report the condition and trend of inshore coral reefs in relation to the extent, frequency and intensity of acute and chronic impacts
- Monitor, assess and report the condition and trend of inshore seagrass meadows in relation to the extent, frequency and intensity of acute and chronic impacts.

Since the 2015-2016 water year, the Reef Water Quality Report Card marine result has been based on averaging the scores for water quality from the [eReefs](#) model output (Robillot *et al.* 2018) with scores for coral and seagrass condition from the Marine Monitoring Program.

The inshore water quality component of the Marine Monitoring Program provides data on physico-chemical water quality parameters including nutrients and sediment concentrations in four Natural Resource Management (NRM) regions, and wet season flood plume exposure and risk to marine communities. Details are not provided in this report and can be found in the [annual technical report for inshore water quality](#) (Waterhouse *et al.* 2021).

## Seagrass condition

77% of seagrass meadows are in the inshore water body. They are fundamental to fisheries productivity and the main food source of dugongs and turtles. Monitoring was conducted at 69 sites at 31 locations in six regions during 2019–2020 by three field teams ([McKenzie et al. 2021](#)) (see **Error! Reference source not found.**). Five major seagrass habitat types (estuarine, coastal intertidal, coastal subtidal, reef intertidal and reef subtidal) are assessed where possible.



*Figure 1. Marine Monitoring Program seagrass survey locations (including Reef Joint Field Management Team and Seagrass-Watch). Not all sites are surveyed every year. Source: (McKenzie et al. 2021).*

Sampling was undertaken at 14 coastal, four estuarine and 12 reef locations (i.e. two or three sites at each location). Reef intertidal sites in the Burdekin and Wet Tropics were paired with a subtidal site. At each location, with the exception of subtidal sites, sampling included two sites nested within 500m of each other. Subtidal sites were not always replicated within locations. Intertidal sites were defined as a 5.5ha area within a relatively homogenous section of a representative seagrass community/meadow. Monitoring occurred in the late dry season (September-November 2019) and late wet season (March/April 2020).

Three indicators were assessed:

- Seagrass abundance (per cent cover) is an assessment of the average per cent cover of seagrass at a monitoring site in relation to the Seagrass Abundance Guidelines (McKenzie 2009).
- Reproductive effort is the ratio of the average number of reproductive structures (spathes, fruits, female and male flowers) of plants on an area basis relative to the long-term average, and provides an indication of the capacity for meadow recovery following disturbances.
- Tissue nutrient composition is an indicator of nutrient enrichment relative to light available for growth (McKenzie *et al.* 2021).

Additional indicators of seagrass condition and resilience include species composition, relative meadow extent and density of seeds in the seed bank (McKenzie *et al.* 2021).

Environmental pressures are also recorded including within-canopy water temperature, within-canopy benthic light, sediment composition as well as macroalgae and epiphyte abundance.

- Within-canopy benthic light is compared to long-term recorded light levels at individual sites as well as daily light thresholds likely to support long-term growth requirements of the species in these habitats (Collier *et al.* 2016).
- Within-canopy temperature is considered in context of the number of days above 35°C. Growth reduction can occur in some species from prolonged warm water exposure (Collier *et al.* 2011; Collier *et al.* 2016). The critical canopy temperature threshold for photoinhibition and acute temperature stress for seagrass is 40°C (Campbell *et al.* 2006).
- Changes in sediment composition can be an indicator of broader environmental changes (such as sediment and organic matter loads and risk of anoxia) and an early-warning indicator of changing species composition.

Additional data on climate and water quality is obtained from the Bureau of Meteorology and from the Marine Monitoring Program inshore water quality component (Waterhouse *et al.* 2021).

## Coral reef condition

Coral reefs comprise 7% of total area of the Great Barrier Reef Marine Park, 3.6% of coral reefs are in the inshore water body.

Monitoring of inshore coral reef communities occurs routinely in the dry season at reefs adjacent to four regions: Wet Tropics, Burdekin, Mackay Whitsunday and Fitzroy (see Figure 2). No reefs are included in Cape York due to logistic and occupational health and safety issues relating to diving in coastal waters in this region.

31 reefs are monitored biennially at two depths under the program, with an additional eight inshore reefs monitored at single depths under the Australian Institute of Marine Science – Long Term Monitoring Program. All are included in the annual assessment of coral condition, although not all reefs are sampled every year (Thompson *et al.* 2021).




Figure 2: Marine Monitoring Program coral survey locations. Reefs were scheduled to be monitored biennially. Purple dots indicate locations monitored as part of the Long Term Monitoring Program conducted by the Australian Institute of Marine Science. Source: (Thompson *et al.* 2021).

Two sites at each reef are permanently marked with fence posts at the beginning of five, 20m-long transects with smaller steel rods at the midpoint and end of each transect. Monitoring is conducted by divers along these transects. They assess community attributes including hard and soft coral cover, the number of hard coral juvenile colonies (up to 5cm in diameter), proportion (per cent) of macroalgae cover, rate of change in coral cover (as an indication of the recovery potential of the reef following a disturbance) and coral community composition (Thompson *et al.* 2021).

## Assessing status against the objectives

### Improved seagrass condition

Three indicators are used to assess and report inshore seagrass condition: abundance, reproductive effort and tissue nutrient status. Further detail about the selection and scoring of these indicators is available in the [annual technical report](#) (McKenzie *et al.* 2021).

The overall grade is the average of the scores of the three indicators for the monitoring year (colour-graded coaster) for the inshore Reef and regions. To calculate the overall score for seagrass, the regional scores were weighted by the relative proportion of World Heritage Area seagrass (shallower than 15m) within that region (see Table 1).

Table 1: Area of seagrass shallower than 15m in each region within the boundaries of the Great Barrier Reef World Heritage Area\*

Region	Area of seagrass (km <sup>2</sup> )	Weighting factor (per cent)
Cape York	2,078	60
Wet Tropics	207	6
Burdekin	587	17
Mackay Whitsunday	215	6
Fitzroy	257	7
Burnett Mary	120	3
World Heritage Area	3,464	100

\* Derived from (McKenzie, Yoshida, Grech *et al.* 2014; McKenzie, Yoshida, and Unsworth 2014; Carter *et al.* 2016; Waterhouse *et al.* 2016).

The online Report Card also shows a graph of the abundance indicator over time (dark blue circles) for the inshore Reef and regions.

### Improved coral condition

Five indicators are used to assess and report on inshore coral reef condition: coral cover, coral cover change, juvenile coral density, coral community composition and proportional macroalgal cover. Further detail about the selection and scoring of these indicators is available in the [annual technical report](#) (Thompson *et al.* 2021).

The overall grade is the average of the scores of the five indicators for the monitoring year (colour-graded coaster) for the inshore Reef and regions. To calculate the overall score for coral, the regional scores were weighted by the relative proportion of the total inshore reef area in the Great Barrier Reef Marine Park that is represented by each of the four monitored regions (see Table 2).

Table 2: Area of inshore reef in each region within the Marine Park\*

Region	Area of inshore reef (km <sup>2</sup> )	Weighting factor (per cent)
Cape York	265	
Wet Tropics	64	20.9
Burdekin	28	9.2
Mackay Whitsunday	117	38.1
Fitzroy	98	31.8
Burnett Mary	5	
Marine Park	577	100

\* Area statistics supplied by the Authority's Spatial Data Centre, 2011

The online Report Card also shows a graph of the coral cover indicator over time (light blue circles) for the inshore Reef and regions.

### Synthesis and integration of data and information

The Reef Water Quality Report Card 2020 provides scores for the condition of inshore water quality, seagrass and coral at Great Barrier Reef-wide and regional scales.

Reef-wide and regional marine scores are unweighted averages of these three indicator scores.

The Marine Monitoring Program provides the coral and seagrass scores, based on [annual technical reports](#) published in the Great Barrier Reef Marine Park Authority's eLibrary.

The eReefs Marine Modelling Program provides the water quality metric for the inshore Reef score based on open coastal waters (Robillot *et al.* 2018).

### Qualitative confidence rankings

A multi-criteria analysis was used to qualitatively score the confidence in each indicator used in the Reef Water Quality Report Card 2020, from low to high. The approach combined expert opinion and direct measures of error for program components where available. Seagrass and coral both received a four-dot confidence ranking (see Figure 3).



Figure 3: Qualitative confidence rankings for seagrass and coral scores. Source: Refer to Appendix A.

## References

- Australia and Queensland governments. 2018. *Reef 2050 Water Quality Improvement Plan 2017-2022*. Brisbane: Queensland Government.  
<https://www.reefplan.qld.gov.au/about/assets/reef-2050-water-quality-improvement-plan-2017-22.pdf>.
- Campbell, S. J., L. J. McKenzie, and S. P. Kerville. 2006. Photosynthetic Responses of Seven Tropical Seagrasses to Elevated Seawater Temperature. *Journal of Experimental Marine Biology and Ecology* 330 (2): 455-468.
- Carter, A. B., S. A. McKenna, M. A. Rasheed, L. McKenzie, and R. Coles. 2016. *Seagrass Mapping Synthesis: A Resource for Coastal Management in the Great Barrier Reef World Heritage Area: Report to the National Science Programme*. Cairns: Reef and Rainforest Research Centre Limited.
- Collier, C. J., MP Adams, L. Langlois, M. Waycott, KR O'Brien, PS Maxwell, and L. McKenzie. 2016. Thresholds for Morphological Response to Light Reduction for Four Tropical Seagrass Species. *Ecological Indicators* 67: 358-366.
- Collier, C. J., S. Uthicke, and M. Waycott. 2011. Thermal Tolerance to Two Seagrass Species at Contrasting Light Levels: Implications for Future Distribution in the Great Barrier Reef. *Limnology and Oceanography* 56 (6): 2200-2210.
- McKenzie, L. J., R. L. Yoshida, A. Grech, and R. Coles. 2014. *Composite of Coastal Seagrass Meadows in Queensland, Australia - November 1984 to June 2010*. PANGAEA.  
<http://doi.pangaea.de/10.1594/PANGAEA.826368>.
- McKenzie, L. J., R. L. Yoshida, and R. K. F. Unsworth. 2014. Disturbance Influences the Invasion of a Seagrass into an Existing Meadow. *Marine Pollution Bulletin* 86 (1-2): 186-196.
- McKenzie, L. J. 2009. *Condition, Trend and Risk in Coastal Habitats: Seagrass Indicators, Distribution and Thresholds of Potential Concern*. Reef and Rainforest Research Centre, Cairns: Marine and Tropical Research Facility. <http://rrrc.org.au/wp-content/uploads/2014/06/113-QDPIF-McKenzie-L-2009-June-Milestone-Report.pdf> ;.
- McKenzie, L. J., C. J. Collier, L. A. Langlois, R. L. Yoshida, J. Uusitalo, and M. Waycott. 2021. *Marine Monitoring Program: Annual Report for Inshore Seagrass Monitoring 2019-20*. Report for the Great Barrier Reef Marine Park Authority. Townsville: Great Barrier Reef Marine Park Authority.  
[http://elibrary.gbrmpa.gov.au/jspui/browse?type=series&value=Marine+Monitoring+Program&sort\\_by=2&order=DESC&rpp=20&etal=0&submit\\_browse=Update](http://elibrary.gbrmpa.gov.au/jspui/browse?type=series&value=Marine+Monitoring+Program&sort_by=2&order=DESC&rpp=20&etal=0&submit_browse=Update;);
- Robillot, C., M. Logan, M. Baird, J. Waterhouse, K. Martin, and B. Schaffelke. 2018. *Testing and Implementation of an Improved Water Quality Index for the 2016 and 2017 Great Barrier Reef Report Cards: Detailed Technical Report*. Cairns: Report to the National Environmental Science Programme. Reef and Rainforest Research Centre Limited.
- Thompson, A., P. Costello, J. Davidson, M. Logan, and G. Coleman. 2021. *Marine Monitoring Program: Annual Report for Inshore Coral Reef Monitoring 2019-20*. Report for the Great Barrier Reef Marine Park Authority. Townsville: Great Barrier Reef Marine Park Authority.  
[http://elibrary.gbrmpa.gov.au/jspui/browse?type=series&value=Marine+Monitoring+Program&sort\\_by=2&order=DESC&rpp=20&etal=0&submit\\_browse=Update](http://elibrary.gbrmpa.gov.au/jspui/browse?type=series&value=Marine+Monitoring+Program&sort_by=2&order=DESC&rpp=20&etal=0&submit_browse=Update).
- Waterhouse, J., J. Brodie, C. Coppo, D. Tracey, E. da Silva, C. Howley, C. Petus, et al. 2016. *Assessment of the Relative Risk of Water Quality to Ecosystems of the Eastern Cape York*

*NRM Region, Great Barrier Reef. A Report to South Cape York Catchments.* Townsville: TropWATER, James Cook University.

Waterhouse, J., R. Gruber, M. Logan, C. Petus, C. Howley, S. Lewis, D. Tracey, et al. 2021. *Marine Monitoring Program: Annual Report for Inshore Water Quality Monitoring 2019-20. Report for the Great Barrier Reef Marine Park Authority.* Townsville: Great Barrier Reef Marine Park Authority.

[http://elibrary.gbrmpa.gov.au/jspui/browse?type=series&value=Marine+Monitoring+Program&sort\\_by=2&order=DESC&rpp=20&etal=0&submit\\_browse=Update](http://elibrary.gbrmpa.gov.au/jspui/browse?type=series&value=Marine+Monitoring+Program&sort_by=2&order=DESC&rpp=20&etal=0&submit_browse=Update).

## Appendix A: Derivation of confidence ranking

A multi-criteria analysis approach was endorsed by the Independent Science Panel in July 2016 and used to qualitatively score the confidence for each key indicator used in the report card. The approach enables the use of expert opinion and measured data.

A multi-criteria analysis identifies the key components that contribute to a problem. These are known as criteria. Each criterion is then scored using a defined set of scoring attributes. The attributes are ranked from those that contribute weakly to the criteria to those that have a strong influence. If the criteria are seen to have different levels of importance for the problem being addressed, they can be weighted accordingly. The strengths of this approach are that it is repeatable, transparent and can include contributions from a range of sources. The weaknesses are that it can be subjective and open to manipulation.

The determination of confidence for the Report Card used five criteria:

- Maturity of methodology (the score is weighted half for these criteria so not to outweigh the importance of the other criteria)
- Validation
- Representativeness
- Directness
- Measured error

### Seagrass

Maturity of methodology (weighting 0.5)	Validation	Representativeness	Directness	Measured error
New or experimental methodology	Survey with no ground truthing	Less than 10% of population survey data	Measurement of data that have conceptual relationship to reported indicator	Error not measured or >25% error
Peer reviewed method	Survey with ground-truthing (not comprehensive)	<b>10%-30% of population survey data</b>	Measurement of data that have a quantifiable relationship to reported indicators	<b>10-25% error</b>
<b>Established methodology in published paper</b>	<b>Survey with extensive on ground validation or directly measured data</b>	30-50% of population	<b>Direct measurement of reported indicator with error</b>	Less than 10% error
3 x 0.5 = 1.5	3	2	3	2

Bolded and grey shading in cells indicates assessment ranking. Total score = 11.5, equates to Four dots.

## Coral

Maturity of methodology (weighting 0.5)	Validation	Representativeness	Directness	Measured error
New or experimental methodology	Survey with no ground truthing	Less than 10% of population survey data	Measurement of data that have conceptual relationship to reported indicator	Error not measured or >25% error
Peer reviewed method	Survey with ground-truthing (not comprehensive)	<b>10%-30% of population survey data</b>	Measurement of data that have a quantifiable relationship to reported indicators	<b>10-25% error</b>
<b>Established methodology in published paper</b>	<b>Survey with extensive on ground validation or directly measured data</b>	30-50% of population	<b>Direct measurement of reported indicator with error</b>	Less than 10% error
3 x 0.5 = 1.5	3	2	3	2

Bolded and grey shading in cells indicates assessment ranking. Total score = 11.5, equates to **Four dots**.

## Glossary

**Ecosystem:** dynamic complex of plant, animal and microorganism communities and their non-living environment interacting as a functional unit.

**Ecosystem health:** ecological processes, biodiversity and function of biological communities is maintained.

**eReefs:** coupled hydrodynamic and biogeochemical models of water quality and ecosystem condition for the Marine Park <<https://research.csiro.au/ereefs/models/>>.

**Guideline value:** a measurable quantity (e.g. concentration) or condition of an indicator for a specific community value below which (or above which, in the case of stressors) there is considered to be a low risk of unacceptable effects occurring to that community value.

**Inshore:** the enclosed coastal and open coastal water bodies combined. These terms are defined and mapped under schedules in the Environmental Protection (Water) Policy.

**Marine Park:** Great Barrier Reef Marine Park.

**Pollutant:** a substance that is present in concentrations that may harm organisms or exceed an environmental quality standard. In this program, the term refers primarily to nutrients, sediment and pesticides.

**Reef 2050 WQIP:** Reef 2050 Water Quality Improvement Plan.

**Reef 2050 Plan:** Reef 2050 Long-Term Sustainability Plan.